

Packed Lunch Standards

Contents

Page 2. Packed Lunch Standards

Page 3. Examples of common foods in each food group which can be included in packed lunches (table 1)

Page. 4 Examples of common foods in sub-groups which should not be included in packed lunches (table 2)

Page 4. Drinks guidance (table 3)

Growing Healthy Families Packed Lunch Standards			
Meal	Essential criteria	Achieved (yes/no)	Comments/suggestions
Lunch	Each day, the following is included in a packed lunch:		
	A portion of starchy carbohydrates (refer to table 1 for examples)		
	A portion of meat, poultry, fish, meat alternative, eggs or pulses (refer to table 1 for examples)		
	A portion of dairy (refer to table 1 for examples)		
	At least one portion of fruit (refer to table 1 for examples)		
	At least one portion of vegetables (refer to table 1 for examples)		
	Water or milk		
	Over the week:		
	Milk or fruit-based desserts only		
	Should include at least three different portions of fruit or vegetables		
	A range of white and wholegrain starchy carbohydrates		
	Limit processed meats, fish and alternatives (refer to table 2 for examples)		
	Avoid packed lunches containing the following foods:		
	Foods high in saturated fat, salt, and/or sugars like cakes, sweet and savoury pastries, biscuits, crisps, chocolate and other confectionary. (refer to table 3 for examples)		
	Drinks other than milk and water (refer to table 3 for examples)		

Table 1. Examples of common foods in each food group which can be included in packed lunches

Starchy carbohydrates	Vegetables fresh/frozen/canned	Fruits fresh/frozen/canned	Meat/meat alternatives	Dairy and alternatives
Bread	Carrots	Bananas	Chicken	Milk
Bagel	Tomatoes	Pears	Lamb	Cheese
Tortilla	Cucumber	Grapes	Pork	Plain yoghurt (full fat)
Wrap	Celery	Kiwi fruit	Beef	Plant-based milk alternatives: unsweetened calcium fortified soya, oat, pea, coconut or almond milks.
Pitta bread	Peppers	Oranges	White fish	
Chapatti	Sweetcorn	Plums	Tuna	
Potato	Mixed salad	Berries	Cod	Milk based desserts
Yam	Green beans	Melon	Haddock	Plain fromage frais
Pasta	Sugar snack peas	Apple	Salmon	Rice pudding
Rice	Broccoli	Pineapple	Sardines	Semolina
Noodles	Cauliflower	Mango	Pulses/Legumes/Lentils	Custard
Couscous		Fruit based desserts	Eggs	Tapioca pudding
Polenta		Fruit/fruit salad (fresh/ tinned)	Quorn/Soya	
		Fruit fool (limit added sugar)	Houmous	
		Fruit crumble (limit added sugar)	Tahini	
		Fruit cake/muffin (sweetened with fresh or dried fruit only)	Nut butters (consult setting allergen policy)	
		Steamed/baked fruit	Tofu	
			Falafel	

Table 2. Examples of common foods in sub-groups which should not be included in packed lunches

Processed meats/ meat alternatives	Foods high in saturated fat, salt and sugar
Corned beef	Chocolate
Pressed meats (Luncheon/Salami/Chicken roll)	Sweets
Sausage (meat/Quorn)	Cakes
Pastry (sausage roll/pasty/meat pie)	Crisps
Crumbed/battered (schnitzel/nuggets/fish fingers/Quorn nuggets/fillets/scotch eggs)	Doughnuts
	Sweet and savoury pastries
	Ice-cream/lollies
	Jelly
	Chocolate mouse

Table 3. Drinks guidance

Suitable			
Water			
Milk	0-1 years	1-2 years	2+ years
	Breastmilk or infant formula	Whole (full fat) or semi-skimmed milk for children growing and eating well	Whole (full fat) or semi-skimmed milk for children growing and eating well
		WHO recommends breastmilk up to 2 years and beyond	
Not suitable			
Skimmed milk or skimmed milk products, these are low in fat and do not provide enough energy for children under 5.			
Fruit juice (even diluted) as these contain large amounts of fruit sugar (fructose) and is acidic, fruit sugar and acid can cause tooth decay in children.			
Fruit juice drinks such as fruit shoots often contain added sugar or sweeteners, and provide little nutritional value.			
Fruit squash, even those labelled 'sugar free' or 'reduced sugar' can contribute to tooth decay and provide little nutritional value.			
Fizzy drinks, even those labelled 'sugar free' or 'reduced sugar' can contribute to tooth decay and provide little nutritional value.			
Flavoured water, even those labelled 'sugar free' or 'reduced sugar' can contribute to tooth decay and provide little nutritional value.			

References

Action for Children, 2017. Voluntary Food and Drink Guidelines for Early Years Settings in England – A Practical Guide. Available at: [Eat Better, Start Better – Foundation Years](#)

Department for Education, 2025. Early Years Foundation Stage Nutrition Guidance. Available at: [*Early Years Foundation Stage nutrition guidance](#)

First Steps Nutrition Trust, 2024. Eating Well: Packed Lunches for 1-4 Year Olds. Available at: [Packed lunches – First Steps Nutrition Trust](#)

First Steps Nutrition Trust, 2018. Good Food Choices and Portion Sizes for 1-4 Year Olds. Available at: [Eating well early years – First Steps Nutrition Trust](#)