

Catering Menu Standards

These catering standards are designed to help early years providers offer nutritious food and drink to children aged 1 to 5 years in their care. In line with EYFS Nutrition Guidance (2025), and Eat Better, Start Better (2017), the guidance supports providers in meeting the EYFS requirement that states: 'Where children are provided with meals, snacks and drinks, these must be healthy, balanced and nutritious'. Following these standards is important to ensure children receive the right amount of nutrients and develop healthy eating habits.

Contents

Page 2. Breakfast and Lunch

Page 3. Lunch and Tea/Light Meal

Page 4. Tea/Light Meal

Page 5. Morning and Afternoon Snack and Entire Menu

Page 6. Entire Menu

Page 7. Examples of common foods in each food group (Table 1)

Page 8. Examples of common foods in sub-groups (Table 2)

Page 9. Drinks (Table 3)









Growing Healthy Families Catering Menu Standards						
Meal	Essential criteria Achieved (yes/no) Comments/suggestions					
	Each day, are the following offered at breakfast:					
	A portion of starchy carbohydrates (cereal/porridge/bread/toast/English muffin/crumpet)					
	A serving of plain milk <u>or</u> other dairy food (yoghurt/cheese)					
	A portion of fruit <u>or</u> vegetables					
5 J	Water					
Breakfast	Over the week, are the following offered at breakfast:					
	Only cereals fortified with vitamins and minerals, labelled low					
	(green) sugar on packaging					
	No high (red) sugar cereals					
	(chocolate flavoured/honey coated/frosted)					
	Offer a variety of wholegrain and white starchy foods each week					
	At least three different starchy/carbohydrate foods					
	(refer to table 1 for examples)	Addisonal				
Meal	Essential criteria	Achieved (yes/no)	Comments/suggestions			
	Each day, are the following offered at lunch:					
	A portion of meat, poultry, fish, meat alternative, eggs or pulses					
	For vegetarian children, a suitable meat alternative similar in					
	texture and appearance to meat-based dish					
	A portion of starchy carbohydrates					
1	(refer to table 1 for examples)					
Lunch	Two different vegetables					
	(refer to table 1 for examples)					
	Water					
	Include a main course and a milk or fruit-based dessert					









	Over the week, are the following offered at lunch:
	Red and white meat at least once
	(refer to table 1 for examples)
	Fish at least once
	(refer to table 1 for examples)
	A variety of wholegrain and white starchy carbohydrates each
	week
	(refer to table 1 for examples)
	Meat alternative or pulses for all children at least once
	(refer to table 1 for examples)
	At least four different vegetarian options such as meat
Lunch	alternatives, eggs and pulses
	Processed meats no more than once
	(sausages/burgers/nuggets/sausage rolls/individual pies/canned
	meat)
	Processed fish no more than once
	(fish fingers/coated fish product)
	Processed meat alternative no more than once
	(vegetarian sausages/burgers/nuggets/pies)
	At least three different starchy carbohydrates
	(refer to table 1 for examples)
	Processed or fried potato products no more than once
	(chips and roast potatoes)
	Tinned pasta products no more than once and choose no added
	sugar option (1)
	(spaghetti hoops)
	No flavoured dried rice, pasta and noodle products
	(packets and pots of instant flavoured noodles/pasta/rice)
	At least three different vegetables
	(refer to table 1 for examples)
	Low sugar and salt baked beans as a vegetable no more than
	Once A finit beautiful and this can be a set of the can be a set o
	A fruit-based dessert at least twice









	A milk-based dessert at least three times					
Meal	Essential criteria	Achieved (yes/no)	Comments/suggestions			
	Each day, are the following offered at tea:					
	A portion of meat, poultry, fish, meat alternative, eggs or pulses					
	For vegetarian children, a suitable meat alternative similar in					
	texture and appearance to meat-based dish					
	A portion of starchy carbohydrates					
	(refer to table 1 for examples)					
	A portion of vegetables or fruit					
	(refer to table 1 for examples)					
	Water					
	Include a main course and a milk or fruit-based dessert					
	Over the week, are the following offered at tea:					
	Red and white meat at least once					
	(refer to table 1 for examples)					
	Fish at least once					
	(refer to table 1 for examples)					
	A variety of wholegrain and white starchy carbohydrates each					
	week					
	(refer to table 1 for examples)					
Tea/Light Meal	Meat alternative or pulses for all children at least once (refer to table 1 for examples)					
Tear Light Mean	At least three different vegetarian options such as meat					
	alternatives, eggs and pulses					
	Processed meats no more than once					
	(sausages/burgers/nuggets/sausage rolls/individual pies/canned					
	meat)					
	Processed fish no more than once					
	(fish fingers/coated fish product)					
	Processed meat alternative no more than once					
	(vegetarian sausages/burgers/nuggets/pies)					
	At least three different starchy carbohydrates					









Meal	Essential criteria	Achieved (yes/no)	Comments/suggestions
	No salty snacks such as crisps		
	No Flavoured dried rice, pasta, and noodle products		
	Cheese or plain yoghurt no more than twice per week		
	or twice a week		
	Meat, fish, eggs, and meat alternatives offered no more than once		
	Over the week, are the following offered at snack time:	<u> </u>	
Snacks	Plain milk and/or water		
Morning and Afternoon	Snack foods are sugar-free or savoury		
	(refer to table 1 for examples)		
	each day		
	Starchy carbohydrate foods offered as part of at least one snack		
	(refer to table 1 for examples, no dried fruit)		
	At least one portion of fruit, and/or vegetable offered every day		
	A choice of at least two different snack each snack time		
	Each day, are the following offered at snacks:	_(,,,,,	
Meal	Essential criteria	Achieved (yes/no)	Comments/suggestions
	A fruit-based or milk-based dessert at least three times		
	once		
	Low sugar and salt baked beans as a vegetable no more than		
	(refer to table 1 for examples)		
	At least three different vegetables		
	(packets and pots of instant flavoured noodles/pasta/rice)		
	(spaghetti hoops) No flavoured dried rice, pasta and noodle products		
	Tinned pasta products no more than once		
	(chips and roast potatoes)		
	Processed or fried potato products no more than once		
	(refer to table 1 for examples)		









	-	т	
	Provide at least three different varieties of starchy carbohydrates		
	each day, and across each meal and snack every week. These		
	should be a variety of wholegrain and white starchy		
	carbohydrates.		
	Access to drinking water throughout day and children encouraged		
	to help themselves		
	Oily fish is provided (such as salmon, sardines, pilchards or		
	mackerel) at least once every three weeks as part of lunch or tea,		
	but not more than twice each week, as it can contain low levels of		
	pollutants. When oily fish is not provided in a main meal, you		
	could try to provide as a snack.		
	Whole (full fat) milk given to children aged from one to two years		
	as their main drink.		
	Children over the age of two can have semi-skimmed milk if they		
	are growing well and eating a healthy, balanced diet.		
	Avoid skimmed milk as this is low in fat and does not provide		
	enough energy for children under five.		
	Choose bread and bread products with lower salt content, those		
	labelled as low (green) or medium (amber).		
Entire Menu	Fat spreads made from vegetable oils, which are high in		
	unsaturated fats such as olive oil used.		
	Use as little oil in cooking as possible, and choose a cooking oil		
	high in unsaturated fats, such as vegetable, rapeseed, sunflower		
	or olive oil.		
	Grill or bake food instead of frying.		
	Limit use of ready-made sauces, condiments, soups, stocks and		
	gravy granules.		
	Use of pastry limited to once a week		
	Avoid foods high in saturated fat, salt and/or sugars (HFSS) such		
	as cakes, biscuits, crisps, chocolate and other confectionary.		
	Limit dairy ice cream and cream to once a week at mealtimes with		
	fruit-based desserts.		









Avoid foods and drinks containing artificial sweeteners. They may
encourage children to prefer overly sweet foods
Do not add salt in cooking
Provide a variety of fruit and vegetables
No processed desserts more than once e.g. angel delight, jelly
cubes, pre-made cheesecake
Do not use tinned fruit, vegetables or pulses with added sugar,
syrup or salt.
No additives or trans-fats used

Table 1. Examples of common foods in each food group					
Starchy carbohydrates	Vegetables fresh/frozen/canned	Fruits fresh/frozen/canned	Meat/meat alternatives	Dairy and alternatives	High Fat, Salt, Sugar (HFSS) food and drinks
Potato	Carrots	Bananas	Beef	Milk	Butter/margarine/oil
Pasta	Green beans	Pears	Lamb	Cheese (Full fat up to 2yrs at least)	Cream/sour cream
White whole grain brown rice	Broccoli	Grapes	Pork	Processed cheese (sticks/triangles/slices)	Crème fraiche
Noodles	Cauliflower	Kiwi fruit	Chicken	Cream cheese (Full fat up to 2yrs at least)	Mascarpone
Couscous	Cabbage	Oranges	White fish	Yoghurt No added sugar (Full fat up to 2yrs at least)	Sugar
Bread/Rolls/English muffins (white/brown/grain)	Courgettes	Plums	Tuna	Fromage frais No added sugar (Full fat up to 2yrs at least)	Honey/syrup/treacle
Tortilla	Peppers	Berries	Cod	Custard	Muffins (cake style)
Pita	Leeks	Melon	Haddock	For children over one year with a milk allergy or with	Biscuits
Chapatti/naan/paratha/roti	Onions	Apple	Pollock	vegan dietary preference,	Cakes
Crackers/crispbread/oatcak e	Okra	Pineapple	Oily fish	unsweetened calcium fortified soya drink can be used as a non-dairy	Doughnuts
Crumpet/pancake/scone	Swede	Mandarin	Mackerel	alternative.	Pastries/Pies
Breakfast cereal	Pak choi	Prunes	Sardines		Ice-cream









Porridge/oats	Peas	Figs	Herrings	Jelly
Semolina	Sweetcorn	Mango	Pilchards	Mousse (e.g. chocolate)
Tapioca	Celery	Dried fruit used in desserts	Salmon	Ice Iollies
Polenta	Cucumber		Eggs	Sweets
	Tomato		Cheese	Chocolate
	Radish		Pulses/Legumes/Lentils	Fizzy drinks
	Beetroot		Quorn/Soya	Squash
	Pulses/lentils served as a		Choose lean cuts of meat	Some canned fruit and
	vegetable alongside meat,		and remove visible fat and	vegetables contain added
	fish or meat alternative.		skin from meat and poultry.	sugar, syrup or salt.

Table 2. Examples of common foods in sub-groups (for menu audit)					
Processed potato products	Processed meats/ meat alternatives	Milk based desserts	Fruit based desserts*	Unsaturated margarine/ cooking oils	
Chips	Corned beef	Rice/Semolina/Tapioca pudding	Fruit/fruit salad (fresh/ tinned)	Vegetable	
Potato waffles	Pressed meats (Luncheon/Salami/Chicken roll)	Milk jelly	Fruit fool	Soya	
Alphabites™	Sausage (meat/Quorn)	Angel delight	Fruit compote	Rapeseed	
Smiles™	Pastry (sausage roll/pasty/meat pie)	Yoghurt	Fruit crumble/pie	Olive	
Potato scone	Crumbed/battered (schnitzel/nuggets/fish fingers/Quorn nuggets/fillets/scotch eggs)	Custard	Eve's pudding	Groundnut/Peanut	
Potato hash		Fromage frais	Fruit cake/muffin (sweetened with fresh or dried fruit)	Sunflower	
Crisps		Cheesecake	Fruit jelly		
			Steamed fruit pudding		









	Baked apple/pear/peach	
	Stewed fruit	
	*fruit based desserts should aim to include 40g fresh fruit or 20g dried fruit per portion	

Table 3. Drin	Table 3. Drinks					
Suitable	Suitable					
Water	Water					
	0-1 years 1-2 years 2+ years					
Milk	Drootmille	Whole (full fat) or semi-skimmed milk for children growing and eating well	Semi-skimmed milk for children growing and eating well			
	Breastmilk WHO recommends breastmilk up to 2 years and beyond					
Not suitable						
Skimmed milk or skimmed milk products, these are low in fat and do not provide enough energy for children under 5.						

Fruit juice (even diluted) as these contain large amounts of fruit sugar (fructose) and is acidic, fruit sugar and acid can cause tooth decay in children.

Fruit juice drinks such as fruit shoots often contain added sugar or sweeteners, and provide little nutritional value.

Fruit squash, even those labelled 'sugar free' or 'reduced sugar' can contribute to tooth decay and provide little nutritional value.

Fizzy drinks, even those labelled 'sugar free' or 'reduced sugar' can contribute to tooth decay and provide little nutritional value.

Flavoured water, even those labelled 'sugar free' or 'reduced sugar' can contribute to tooth decay and provide little nutritional value.









References

Action for Children, 2017. Eat Better, Start Better. Voluntary Food and Drink Guidelines for Early Years Settings in England – A Practical Guide. Available at: <u>Eat Better</u>, <u>Start Better</u> – Foundation Years

Department for Education, 2025. Early Years Foundation Stage Nutrition Guidance. Available at: *Early Years Foundation Stage nutrition guidance

First Steps Nutrition Trust., 2024. Eating Well: Packed Lunches for 1-4 Year Olds. Available at: Packed lunches — First Steps Nutrition Trust.

First Steps Nutrition Trust, 2018. Good Food Choices and Portion Sizes for 1-4 Year Olds. Available at: <u>Eating well early years — First Steps</u>

<u>Nutrition Trust</u>







